Appendix I—Soil Characteristics of North Campus

TABLE 1. Classification of major soil series occurring in the North Campus area (USDA Soil Conservation Service and Alaska Agricultural Experiment Station 1963)

காகைக் படங்கள் கடைக்கள் கடைக்கள் கூடன் இறக்கள் தொகைப்படித்தில் இதிக்கள் திருந்துக்கள் கட்சு கட்சு கட்சு காண கா பிருந்து

ngineering properties					Soil Series Rating for e		
ream	Minto				Fairbanks		Goldst
ierie innitatoisis (in stape sa saec	e gregier in Constant e	08/29/Chone 11, finite	tv.o≾er theo 7	rene To Stevere timilations on surpres o 35 stope, steep.nese	necen n	i eo wet, dévéré iba naradia, ar bieorito hav portrairost restricta use	wh <u>g: []]]</u> Waterere (1) aer then 3%, wot, it
es ése to thermokarst pilting Small e		nmoroial buildings	Severe	ilmitations over 7% slope, too steep	,	Yee wet, severe restrictions, subject to Rooding, Severe restriction permatrost restricts use	
' I Jo⊷ Wat, isavaik	a taşan konolea işi kinin dir.	bevere units pitting	ส์ดอย่ะ	×je oʻönde€aon•nj prenge•karst i -	_2): 34, pr	≫ās anaistreets – † †i rrost ∞ta∾n .ikaņ	1
7% Too wet, ævere restrictions		Moderate i	Moderate limitations due to wetness		Picnic areas Moderate lin siope, too str		ions on siopes greater than
2%	Initations at 86665 g	Minderate eaterthan 7% soo stee	roitatiq P	rs ive 's we'ress swere	Pigyarq sibpe"so	wnós. Mosierate limitat o steep	ខែឧត ខ្លា <u>នខ្លែកទុស</u> លាកលាកដាក់ព្រះ រ 1
Restrictions due to the mokarst pitting, erosion, and steepness			1 .	Pond reservoir area	- Piping likely		Permafrost restricts use
gth, piping likely Soil has low strength or bearing strength, compacts poorly, susceptible to pining				Embankments, dikes, levees	Soil has l piping Ji	Soil has low bearing strength, hard to compact, Soil has low piping Jikely	

Thermokarst mounds in the Fairbanks area are polygonal or circular hummocks 10 to 50 feet in diameter, 1-8 feet in height and are composed of loess. They are commonest in cultivated field but there are also some in abandoned fields now reforested. In some fields the mounds are separated by trenches 1-5 feet wide, but in others the trenches are poorly developed. The trenches form as a result of melting of ice masses.

A field on a gentle north-facing slope at the AES has the best developed mounds and the most detailed record in the Fairbanks area. The surface of the field was smooth before clearing in 1908. By 1922 pronounced individual and connected depressions had formed and by 1926 some trenches between mounds were as much as 5 feet deep. Cultivation stopped a year or two later because the irregular topography formed by the pits and mounds was dangerous to the operation of farm machinery. The field then was seeded to pasture. By 1938, the mounds were 3 to 8 feet high and about 20 to 50 feet in diameter. Rockie studied the field in November 1938 in order to determine whether the ice still was actively melting "A tractor bulldozer was used to remove the upper part of every hummock and fill each pit until the land surface assumed approximately a uniform slope. The surface remained smooth for nearly a year but in July 1939 irregularities began to form. In succeeding years polygonal mounds formed as the ground surface subsided over melting ice.

Mounds in the test area smoothed in 1938 were as large and as high as those in the part of the field that had not been smoothed when the writer first studied the field in 1947. Maximum mound height was 8 feet. Comparisons of aerial photos with those taken 10 years earlier reveal that in 1948 mounds were about the same size and shape and in the same position as in 1938. Probing with a soil auger on 14 July 1948, revealed no ice or frozen ground at a depth of 9 f

October 22, 2004 by _____